

**Rescue Dislodged undilated iliac stent at iliac bifurcation successfully**

Wenjung Chung

<sup>1</sup>Department of Internal Medicine, Division of Cardiology,, Kaohsiung Chang Gung memorial hospital, Taiwan, R.O.C.

63 y/o female with history of DM,ESRD on HD,HTN admitted for Rt foot chronic wound. Via LT CFA access(6Fr Cook ANL2 55cm), we complete Rt below knee artery EVT with balloon angioplasty and DES to Rt m-SFA.

For Rt CIA stenotic lesion with pressure gradient, we deliver a balloon expandable stent(Vis-pro 8.0x27) to Rt CIA after initial balloon pre-dilation but stent tip stuck stuck by Rt CIA lesion. When we removed the undilated stent from sheath, the stent dislodged at Lt CIA and stent balloon was removed from sheath. Undilated stent balloon failed to pass the dislodged stent. So,we used a new Mustang 8.0x40mm to inflate the distal stent edge to create a Funnel-shape opening stent. So,we easily delivered a smaller balloon (Mustang 5.0x20mm ) to pass the dislodged stent and relocated at Lt ostial CIA. Then, we deployed the dislodged stent at LT ostial CIA after balloon angioplasty.

Finally, we advanced the sheath (Cook ANL2 55cm) to Rt iliac artery passing the initial Rt CIA lesion. We deployed additional iliac stent to (Express LD 8.0x27) at Rt CIA ostium successfully.